



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386

- Home
- University
- Faculties
- Institutions
- Studying
- Research
- International Affairs
- Excellence Initiative

- Alumni
- Employees
- Friends & Supporters
- Press & News
- Business & Industry

SEARCH Extended search

Web pages People Library Courses




Contact | Deutsch

Home > Press > Press Releases >

Elucidating Environmental History with 100 Million Laser Beams

Press Release No. 82/2013
29 April 2013

Laser scanning and subsurface geodata fused for 3D reconstruction of karst depressions on Crete



Video https://www.youtube.com/watch?v=_9jgPC6zGf8

By combining high-resolution surface data obtained from laser scanning with subsurface geodata, scientists from Heidelberg University have succeeded for the first time in providing a full picture of so-called karst depressions on the island of Crete, including a three-dimensional view into the subsurface structure of these funnel-shaped hollows. This new 3D representation method has been developed under the leadership of junior professor Dr. Bernhard Höfle at Heidelberg University's Institute of

Geography. It is ideal for in-depth analyses at the interface between geosciences and ancient studies. The sediment infills of karst depressions provide terrestrial archives of great value for the reconstruction of environmental scenarios from the past.

For thousands of years, karst landforms and particularly karst depressions like dolines, for example, have been important sites of human husbandry. There is evidence from as far back as the 2nd century BC that such depressions were used for agriculture and livestock breeding, including in the mountainous regions of Crete. Due to their funnel-shaped form, dolines serve as "material traps" in which loose sediments, archaeological finds or volcanic ashes can accumulate. "These infills can supply important information on former climatic conditions, vegetation structure and also on human impact through land use", says Dr. Christoph Siart, a fellow researcher of Prof. Höfle's.

So far, say the Heidelberg geographers, karst depressions have usually only been examined in connection with drilling of sediment cores. These provide insights into the subsurface structure, albeit of a discrete nature, and have been drawn upon in conjunction with geomorphological surface finds to propose explanations for the genesis and function of dolines. With the aid of the new 3D data-modelling method developed in Heidelberg, the scientists have now succeeded in combining the two-dimensional subsurface data with high-resolution 3D surface data. To accomplish this, surface topography data were acquired with the aid of terrestrial laser scanning. Two-dimensional views of various cross-sections of the subsurface of the sediment-filled dolines were achieved with a combination of various geophysical measuring procedures.

The fusion of these data now makes it feasible to undertake soundly substantiated geomorphometric analyses. "For example, we can determine the volume or undertake a digital measurement of the depressions in a virtual three-dimensional model", says Prof. Höfle. "That means we have created the basis for first-ever statements on the genesis, the sediment infill process and the age of the dolines. This is of immense significance for the reconstruction of the environmental history because it supplies a holistic view of geomorphological forms via a combination of surface and subsurface data and thus helps to understand the local processes that ultimately led to the formation of the landscape as we know it today."

Data collection and methodological development took place in the framework of the projects "Reconstruction of Holocene Environmental Change on Crete" and "Geoinformatics and 3D Geoinformation Technology" conducted in the physical geography and geoinformatics research groups of Heidelberg University's Institute of Geography. For more information, go to <http://gis.science.uni-hd.de>.

Note for news desks

Digital pictures are available from the press office.

Original publication

C. Siart, M. Forbriger, E. Nowaczinski, S. Hecht & B. Höfle: Fusion of multi-resolution surface (terrestrial laser scanning) and subsurface geodata (ERT, SRT) for karst landform investigation and geomorphometric quantification; *Earth Surface Processes and Landforms* (2013), doi: 10.1002/esp.3394

Contact

Junior professor Dr. Bernhard Höfle
Institute of Geography and Heidelberg Center for the Environment (HCE)
Phone: +49 6221 54-5594
hoeffle@uni-heidelberg.de

Communications and Marketing
Press Office
phone: +49 6221 54-2311
presse@rektorat.uni-heidelberg.de

Editor: Email

